

BINARY PULSARS AND RELATIVISTIC GRAVITY

Nobel Lecture, December 8, 1993

by

JOSEPH H. TAYLOR, JR.

Princeton University, Department of Physics, Princeton, NJ 08544, USA

I. SEARCH AND DISCOVERY

Work leading to the discovery of the first pulsar in a binary system began more than twenty years ago, so it seems reasonable to begin with a bit of history. Pulsars burst onto the scene (1) in February 1968, about a month after I completed my PhD at Harvard University. Having accepted an offer to remain there on a post-doctoral fellowship, I was looking for an interesting new project in radio astronomy. When *Nature* announced the discovery of a strange new rapidly pulsating radio source, I immediately drafted a proposal, together with Harvard colleagues, to observe it with the 92 m radio telescope of the National Radio Astronomy Observatory. By late spring we had detected and studied all four of the pulsars which by then had been discovered by the Cambridge group, and I began thinking about how to find further examples of these fascinating objects, which were already thought likely to be neutron stars. Pulsar signals are generally quite weak, but have some unique characteristics that suggest effective search strategies. Their otherwise noise-like signals are modulated by periodic, impulsive waveforms; as a consequence, dispersive propagation through the interstellar medium makes the narrow pulses appear to sweep rapidly downward in frequency. I devised a computer algorithm for recognizing such periodic, dispersed signals in the inevitable background noise, and in June 1968 we used it to discover the fifth known pulsar (2).

Since pulsar emissions exhibited a wide variety of new and unexpected phenomena, we observers put considerable effort into recording and studying their details and peculiarities. A pulsar model based on strongly magnetized, rapidly spinning neutron stars was soon established as consistent with most of the known facts (3). The model was strongly supported by the discovery of pulsars inside the glowing, gaseous remnants of two supernova explosions, where neutron stars should be created (4, 5), and also by an observed gradual lengthening of pulsar periods (6) and polarization measurements that clearly suggested a rotating source (7). The electro-dynamical properties of a spinning, magnetized neutron star were studied theoretically (8) and shown to be plausibly capable of generating broadband radio noise

detectable over interstellar distances. However, the rich diversity of the observed radio pulses suggested magnetospheric complexities far beyond those readily incorporated in theoretical models. Many of us suspected that detailed understanding of the pulsar emission mechanism might be a long time coming - and that, in any case, the details might not turn out to be fundamentally illuminating.

In September 1969, I joined the faculty at the University of Massachusetts, where a small group of us planned to build a large, cheap radio telescope especially for observing pulsars. Our telescope took several years to build, and during this time it became clear that whatever the significance of their magnetospheric physics, pulsars were interesting and potentially important to study for quite different reasons. As the collapsed remnants of supernova explosions, they could provide unique experimental data on the final stages of stellar evolution, as well as an opportunity to study the properties of nuclear matter in bulk. Moreover, many pulsars had been shown to be remarkably stable natural clocks (9), thus providing an alluring challenge to the experimenter, with consequences and applications about which we could only speculate at the time. For such reasons as these, by the summer of 1972 I was devoting a large portion of my research time to the pursuit of accurate timing measurements of known pulsars, using our new telescope in western Massachusetts, and to planning a large-scale pulsar search that would use bigger telescopes at the national facilities.

I suspect it is not unusual for an experiment's motivation to depend, at least in part, on private thoughts quite unrelated to avowed scientific goals. The challenge of a good intellectual puzzle, and the quiet satisfaction of finding a clever solution, must certainly rank highly among my own incentives and rewards. If an experiment seems difficult to do, but plausibly has interesting consequences, one feels compelled to give it a try. Pulsar searching is the perfect example: it's clear that there must be lots of pulsars out there, and once identified, they are not so very hard to observe. But finding each one for the first time is a formidable task, one that can become a sort of detective game. To play the game you invent an efficient way of gathering clues, sorting, and assessing them, hoping to discover the identities and celestial locations of all the guilty parties.

Most of the several dozen pulsars known in early 1972 were discovered by examination of strip-chart records, without benefit of further signal processing. Nevertheless, it was clear that digital computer techniques would be essential parts of more sensitive surveys. Detecting new pulsars is necessarily a multi-dimensional process; in addition to the usual variables of two spatial coordinates, one must also search thoroughly over wide ranges of period and dispersion measure. Our first pulsar survey, in 1968, sought evidence of pulsar signals by computing the discrete Fourier transforms of long sequences of intensity samples, allowing for the expected narrow pulse shapes by summing the amplitudes of a dozen or more harmonically related frequency components. I first described this basic algorithm (10) as part of a discussion of pulsar search techniques, in 1969. An efficient dispersion-

compensating algorithm was conceived and implemented soon afterward (11, 12), permitting extension of the method to two dimensions. Computerized searches over period and dispersion measure, using these basic algorithms, have by now accounted for discovery of the vast majority of nearly 600 known pulsars, including forty in binary systems (13, 14).

In addition to private stimuli related to “the thrill of the chase,” my outwardly expressed scientific motivation for planning an extensive pulsar survey in 1972 was a desire to double or triple the number of known pulsars. I had in mind the need for a more solid statistical basis for drawing conclusions about the total number of pulsars in the Galaxy, their spatial distribution, how they fit into the scheme of stellar evolution, and so on. I also realized (15) that it would be highly desirable “. . . to find even *one* example of a pulsar in a binary system, for measurement of its parameters could yield the pulsar mass, an extremely important number.” Little did I suspect that just such a discovery would be made, or that it would have much greater significance that anyone had foreseen! In addition to its own importance, the binary pulsar PSR 1913 + 16 is now recognized as the harbinger of a new class of unusually short-period pulsars with numerous important applications.

An up-to-date map of known pulsars on the celestial sphere is shown in Figure 1. The binary pulsar PSR 1913 + 16 is found in a clump of objects close to the Galactic plane around longitude 50° , a part of the sky that passes directly overhead at the latitude of Puerto Rico. Forty of these pulsars, including PSR 1913 + 16, were discovered in the survey that Russell Hulse and I carried out with the 305 m Arecibo telescope (16 - 18). Figure 2 illustrates the periods and spin-down rates of known pulsars, with those in binary systems marked by larger circles around the dots. All radio pulsars slow down gradually in their own rest frames, but the slowdown rates vary




Fig. 1: Distribution of 558 pulsars in Galactic coordinates. The Galactic center is in the middle, and longitude increases to the left.




Fig. 2: Periods and period derivatives of known pulsars. Binary pulsars, denoted by larger circles around the dots, generally have short periods and small derivatives. Symbols aligned near the bottom represent pulsars for which the slow-down rate has not yet been measured.

over nine orders of magnitude. Figure 2 makes it clear that binary pulsars are special in this regard. With few exceptions, they have unusually small values of both period and period derivative - an important fact which helps to make them especially suitable for high-precision timing measurements.

Much of the detailed implementation and execution of our 1973 - 74 Arecibo survey was carried out by Russell Hulse. He describes that work, and particularly the discovery of PSR 1913 + 16, in his accompanying lecture (19). The significant consequences of our discovery have required accurate timing measurements extending over many years, and since

1974-76 I have pursued these with a number of other collaborators. I shall now turn to a description of these observations.

II. CLOCK-COMPARISON EXPERIMENTS

Pulsar timing experiments are straightforward in concept: one measures pulse times of arrival (TOAs) at the telescope, and compares them with time kept by a stable reference clock. A remarkable wealth of information about a pulsar's spin, location in space, and orbital motion can be obtained from such simple measurements. For binary pulsars, especially, the task of analyzing a sequence of TOAs often assumes the guise of another intricate detective game. Principal clues in this game are the recorded TOAs. The first and most difficult objective is the assignment of unambiguous pulse numbers to each TOA, despite the fact that some of the observations may be separated by months or even years from their nearest neighbors. During such inevitable gaps in the data, a pulsar may have rotated through as many as 10^7 - 10^{10} turns, and in order to extract the maximum information content from the data, these integers must be recovered exactly. Fortunately, the correct sequence of pulse numbers is easily recognized, once attained, so you can tell when the game has been "won."

A block diagram of equipment used for recent pulsar timing observations (20) at Arecibo is shown in Figure 3. Incoming radio-frequency signals from the antenna are amplified, converted to intermediate frequency, and passed




Fig. 3: Simplified block diagram of equipment used for timing pulsars at Arecibo.

through a multichannel spectrometer equipped with square-law detectors. A bank of digital signal averagers accumulates estimates of a pulsar's periodic waveform in each spectral channel, using a pre-computed digital ephemeris and circuitry synchronized with the observatory's master clock. A programmable synthesizer, its output frequency adjusted once a second in a phase-continuous manner, compensates for changing Doppler shifts caused by accelerations of the pulsar and the telescope. Average profiles are recorded once every few minutes, together with appropriate time tags. A




Fig. 4: Pulse profiles obtained on April 24, 1992 during a five-minute observation of PSR 1913 + 16. The characteristic double-peaked shape, clearly seen in the de-dispersed profile at the bottom, is also discernible in the 32 individual spectral channels.

log is kept of small measured offsets (typically of order $1\mu\text{s}$) between the observatory clock and the best available standards at national time-keeping laboratories, with time transfer accomplished via satellites in the Global Positioning System.

An example of pulse profiles recorded during timing observations of PSR 1913 + 16 is presented in Figure 4, which shows intensity profiles for 32 spectral channels spanning the frequency range 1383 - 1423 MHz, followed by a “de-dispersed” profile at the bottom. In a five-minute observation such as this, the signal-to-noise ratio is just high enough for the double-peaked pulse shape of PSR 1913 + 16 to be evident in the individual channels. Pulse arrival times are determined by measuring the phase offset between each observed profile and a long-term average with much higher signal-to-noise ratio. Differential dispersive delays are removed, the adjusted offsets are averaged over all channels, and the resulting mean value is added to the time-tag to obtain an equivalent TOA. Nearly 5,000 such five-minute measurements have been obtained for PSR 1913 + 16 since 1974, using essentially this technique. Through a number of improvements in the data-taking systems (21-26), the typical uncertainties have been reduced from around $300\mu\text{s}$ in 1974 to $15\text{-}20\text{-}\mu\text{s}$ since 1981.

III. MODEL FITTING

In the process of data analysis, each measured topocentric TOA, say t_{obs} , must be transformed to a corresponding proper time of emission T in the pulsar frame. Under the assumption of a deterministic spin-down law, the rotational phase of the pulsar is given by

$$\phi(T) = \nu T + \frac{1}{\alpha} \dot{\nu} T^2, \quad (1)$$

where ϕ is measured in cycles, $\nu \equiv 1/P$ is the rotation frequency, P the period, $\dot{\nu}$ the slowdown rate. Since a topocentric TOA is a relativistic space-time event, it must be transformed as a four-vector. The telescope’s location at the time of a measurement is obtained from a numerically integrated solar-system model, together with published data on the Earth’s unpredictable rotational variations. As a first step one normally transforms to the solar-system barycenter, using the weak-field, slow-motion limit of general relativity. The necessary equations include terms depending on the positions, velocities, and masses of all significant solar-system bodies. Next, one accounts for propagation effects in the interstellar medium; and finally, for the orbital motion of the pulsar itself.

With presently achievable accuracies, all significant terms in the relativistic transformation can be summarized in the single equation

$$T = t_{\text{obs}} - t_0 + \Delta_{\text{C}} - D/f^2 + \Delta_{\text{R}\odot}(\alpha, \delta, \mu_{\alpha}, \mu_{\delta}, \pi) + \Delta_{\text{E}\odot} - \Delta_{\text{S}\odot}(\alpha, \delta) - \Delta_{\text{R}}(\mathbf{x}, e, P_b, T_0, \omega, \dot{\omega}, \dot{P}_b) - \Delta_{\text{E}}(\gamma) - \Delta_{\text{S}}(r, s). \quad (2)$$

Here t_0 is a nominal equivalent TOA at the solar system barycenter;

Δ_C represents measured clock offsets; D/f^2 is the dispersive delay for propagation at frequency f through the interstellar medium; $\Delta_{R\odot}$, $\Delta_{E\odot}$, and $\Delta_{S\odot}$ are propagation delays and relativistic time adjustments within the solar system; and Δ_R , Δ_E , and Δ_s are similar terms for effects within a binary pulsar's orbit. Subscripts on the various Δ 's indicate the nature of the time-dependent delays, which include "Römer," "Einstein," and "Shapiro" delays in the solar system and in the pulsar orbit. The Römer terms have amplitudes comparable to the orbital periods times $v/2\pi I c$, where v is the orbital velocity and c the speed of light. The Einstein terms, representing the integrated effects of gravitational redshift and time dilation, are smaller by another factor ev/c , where e is the orbital eccentricity. The Shapiro time delay is a result of reduced velocities that accompany the well-known bending of light rays propagating close to a massive object. The delay amounts to about $120 \mu\text{s}$ for one-way lines of sight grazing the Sun, and the magnitude depends logarithmically on the angular impact parameter. The corresponding delay within a binary pulsar orbit depends on the companion star's mass, the orbital phase, and the inclination i between the orbital angular momentum and the line of sight.

Figure 5 illustrates the combined orbital delay $\Delta_R + \Delta_E + \Delta_s$ for PSR 1913 + 16, plotted as a function of orbital phase. Despite the fact that the Einstein and Shapiro effects are orders of magnitude smaller than the Römer delay, they can still be measured separately if the precision of available TOAs is high enough. In fact, the available precision is very high




Fig. 5: Orbital delays observed for PSR 1913 + 16 during July, 1988. The uncertainty of an individual five-minute measurement is typically 50,000 times smaller than the error bar shown.

indeed, as one can see from the lone data point shown in Figure 5 with $50,000\sigma$ error bars.

Eqs. (1) and (2) have been written to show explicitly the most significant dependences of pulsar phase on as many as nineteen *a priori* unknowns. In addition to the rotational frequency ν and spin-down rate it, these phenomenological parameters include a reference arrival time t_0 , the dispersion constant D , celestial coordinates α and δ , proper-motion terms μ_α and μ_δ , and annual parallax a . For binary pulsars the terms on the second line of Eq. (2), with as many as ten significant orbital parameters, are also required. The additional parameters include five that would be necessary even in a purely Keplerian analysis of orbital motion: the projected semi-major axis $x \equiv a_1 \sin i/c$, eccentricity e , binary period P_b , longitude of periastron ω , and time of periastron T_0 . If the experimental precision is high enough, relativistic effects can yield the values of five further "post-Keplerian" parameters: the secular derivatives $\dot{\omega}$ and \dot{P}_b , the Einstein parameter γ , and the range and shape of the orbital Shapiro delay, r and $s \equiv \sin i$. Several earlier versions of this formalism for treating timing measurements of binary pulsars exist (27-29), and have been historically important to our progress with the PSR 1913 + 16 experiment. The elegant framework outlined here was derived during 1985-86 by Damour and Deruelle (30,31).

Model parameters are extracted from a set of TOAs by calculating the pulsar phases $\phi(T)$ from Eq. (1) and minimizing the weighted sum of squared residuals,

$$\chi^2 = \sum_{i=1}^N \left(\frac{\phi(T_i) - n_i}{\sigma_i/P} \right)^2, \quad (3)$$

with respect to each parameter to be determined. In this equation, n_i is the closest integer to $\phi(T_i)$, and σ_i is the estimated uncertainty of the i 'th TOA. In a valid and reliable solution the value of χ^2 will be close to the number of degrees of freedom, i.e., the number of measurements N minus the number of adjustable parameters. Parameter errors so large that the closest integer to $\phi(T_i)$ may not be the correct pulse number are invariably accompanied by huge increases in χ^2 ; this is the reason for my earlier statement that correct pulse numbering is easily recognizable, once attained. In addition to providing a list of fitted parameter values and their estimated uncertainties, the least-squares solution produces a set of post-fit residuals, or differences between measured TOAs and those predicted by the model (see Figure 6). The post-fit residuals are carefully examined for evidence of systematic trends that might suggest experimental errors, or some inadequacy in the astrophysical model, or perhaps deep physical truths about the nature of gravity.

Necessarily, some model parameters will be easier to measure than others. When many TOAs are available, spaced over many months or years, it generally follows that at least the pulsar's celestial coordinates, spin parameters, and Keplerian orbital elements will be measurable with high precision, often as many as 6-14 significant digits. As we will see, the relativistic




Fig. 6: Schematic diagram of the analysis of pulsar timing measurements carried out by the computer program TEMPO. The essential functions are all described in the text.

parameters of binary pulsar orbits are generally much more difficult to measure - but the potential rewards for doing so are substantial.

IV. THE NEWTONIAN LIMIT

Thirty-five binary pulsar systems have now been studied well enough to determine their basic parameters, including the Keplerian orbital elements, with good accuracy. For each system the orbital period P_b and projected semi-major axis x can be combined to give the mass function,

$$f_1(m_1, m_2, s) = \frac{(m_2 s)^3}{(m_1 + m_2)^2} = \frac{x^3}{T_\odot (P_b/2\pi)^2}. \quad (4)$$

Here m_1 and m_2 are the masses of the pulsar and companion in units of the Sun's mass, M_\odot ; I use the shorthand notations $s \equiv \sin i$, $T_\odot \equiv GM_\odot/c^3 = 4.925490947 \cdot 10^6$ s, where G is the Newtonian constant of gravity. In the absence of other information, the mass function cannot provide unique solutions for m_1 , m_2 , or s . Nevertheless, likely values of m_2 can be estimated by assuming a pulsar mass close to $1.4 M_\odot$ (the Chandrasekhar limit for white dwarfs) and the median value $\cos i = 0.5$, which implies $s = 0.87$. With this approach one can distinguish three categories of binary pulsars, which I




Fig. 7: Masses of the companions of binary pulsars, plotted as a function of orbital eccentricity. Near the marked location of PSR 1913 + 16, three distinct symbols have merged into one; these three binary systems, as well as their two nearest neighbors in the graph, are thought to be pairs of neutron stars. The two pulsars at the upper right are accompanied by high-mass main-sequence stars, while the remainder are believed to have white-dwarf companions.

shall discuss by reference to Figure 7: a plot of binary-pulsar companion masses versus orbital eccentricities.

Twenty-eight of the binary systems in Figure 7 have orbital eccentricities $e < 0.25$ and low-mass companions likely to be degenerate dwarfs. Most of these have nearly circular orbits; indeed, the only ones with eccentricities more than a few percent are located in globular clusters, and their orbits have probably been perturbed by near collisions with other stars. Five of the binaries have much larger eccentricities and likely companion masses of $0.8M_{\odot}$ or more; these systems are thought to be pairs of neutron stars, one of which is the detectable pulsar. Their large orbital eccentricities are almost certainly the result of rapid ejection of mass in the supernova explosion creating the second neutron star. Finally, at the upper right of Figure 7 we find two binary pulsars that move in eccentric orbits around high-mass main-sequence stars. These systems have not yet evolved to the stage of a second supernova explosion. Unlike the binary pulsars with compact companions, these two systems have orbits that could be significantly modified by complications such as tidal forces or mass loss.

V. GENERAL RELATIVITY AS A TOOL

As Russell Hulse and I suggested (17) in the discovery paper for PSR 1913 + 16, it should be possible to combine measurements of relativistic orbital parameters with the mass function, thereby determining masses of both stars and the orbital inclination. In the post-Keplerian (PK) framework outlined above, each measured PK parameter defines a unique curve in the (m_1, m_2) plane, valid within a specified theory of gravity. Experimental values for any two PK parameters (say $\dot{\omega}$ and γ , or perhaps r and s) establish the values of m_1 , m_2 , and s unambiguously. In general relativity the equations for the five most significant PK parameters are as follows (25, 31, 32):

$$\dot{\omega} = 3 \left(\frac{P_b}{2\pi} \right)^{-5/3} (T_{\odot} M)^{2/3} (1 - e^2)^{-1}, \quad (5)$$

$$\gamma = e \left(\frac{P_b}{2\pi} \right)^{1/3} T_{\odot}^{2/3} M^{-4/3} m_2 (m_1 + 2m_2), \quad (6)$$

$$\dot{P}_b = - \frac{192\pi}{5} \left(\frac{P_b}{2\pi} \right)^{-5/3} \left(1 + \frac{73}{24} e^2 + \frac{37}{96} e^4 \right) (1 - e^2)^{-7/2} T_{\odot}^{5/3} m_1 m_2 M^{-1/3}, \quad (7)$$

$$r = T_{\odot} m_2, \quad (8)$$

$$s = x \left(\frac{P_b}{2\pi} \right)^{-2/3} T_{\odot}^{-1/3} M^{2/3} m_2^{-1}. \quad (9)$$

Again the masses m_1 , m_2 , and $M \equiv m_1 + m_2$ are expressed in solar units. I emphasize that the left-hand sides of Eqs. (5-9) represent directly measurable quantities, at least in principle. Any two such measurements, together with the well-determined values of e and P_b , will yield solutions for m_1 and m_2 as well as explicit predictions for the remaining PK parameters.

The binary systems most likely to yield measurable PK parameters are those with large masses and high eccentricities and which are astrophysically "clean," so that their orbits are overwhelmingly dominated by the gravitational interactions between two compact masses. The five pulsars clustered near PSR 1913 + 16 in Figure 7 would seem to be especially good candidates, and this has been borne out in practice. In the most favorable circumstances, even binary pulsars with low-mass companions and nearly circular orbits can yield significant post-Keplerian measurements. The best present example is PSR 1855 + 09: its orbital plane is nearly parallel to the line of sight, greatly magnifying the orbital Shapiro delay. The relevant measurements (33-35) are illustrated in Figure 8, together with the fitted function $\Delta_S(r, s)$, in this case closely approximated by

$$\Delta_S = - 2r \log (1 - s \cos[2\pi(\phi - \phi_0)]) \quad (10)$$

where ϕ is the orbital phase in cycles and $\phi_0 = 0.4823$ the phase of superior conjunction. The fitted values of r and s yield the masses $m_1 = 1.50^{+0.26}_{-0.14}$, m_2




Fig. 8: Measurements of the Shapiro time delay in the PSR 1855+09 system. The theoretical curve corresponds to Eq. (10), and the fitted values of τ and s can be used to determine the masses of the pulsar and companion star.

$= 0.258^{+0.028}_{-0.016}$. In a similar way, all binary pulsars with two measurable PK parameters yield solutions for their component masses. At present, most of the experimental data on the masses of neutron stars (see Figure 9) come from such timing analyses of binary pulsar systems (36, 37, and references therein).

VI. TESTING FOR GRAVITATIONAL WAVES

If three or more post-Keplerian parameters can be measured for a particular pulsar, the system becomes over-determined, and the extra experimental degrees of freedom transform it into a calibrated laboratory for testing relativistic gravity. Each measurable PK parameter beyond the first two provides an explicit, quantitative test. Because the velocities and gravita-




Fig. 9: The masses of ten neutron stars, measured by observing relativistic effects in binary pulsar orbits.

tional energies in a high-mass binary pulsar system can be significantly relativistic, strong-field and radiative effects come into play. Two binary pulsars, PSRs 1913 + 16 and 1534 + 12, have now been timed well enough and long enough to yield three or more PK parameters. Each one provides significant tests of gravitation beyond the weak-field, slow-motion limit (32, 38).

PSR 1913+16 has an orbital period $P_b \approx 7.8$ h, eccentricity $e \approx 0.62$ and mass function $f_1 \approx 0.13 M_{\odot}$. With the available data quality and time span, the Keplerian orbital parameters are actually determined with fractional accuracies of a few parts per million, or better. In addition, the PK parameters $\dot{\omega}$, γ , and b , are determined with fractional accuracies better than $3 \cdot 10^{-6}$, $5 \cdot 10^{-4}$, and $4 \cdot 10^{-3}$, respectively (25, 39). Within any viable relativistic theory of gravity, the values of $\dot{\omega}$ and γ yield the values of m_1 and m_2 and a corresponding prediction for \dot{P}_b arising from the damping effects of gravitational radiation. At present levels of accuracy, a small kinematic correction (approximately 0.5% of the observed \dot{P}_b) must be included to account for accelerations of the solar system and the binary pulsar system in




Fig. 10: Accumulated shift of the times of periastron in the PSR 1913+16 system, relative to an assumed orbit with constant period. The parabolic curve represents the general relativistic prediction for energy losses from gravitational radiation.

the Galactic gravitational field (40). After doing so, we find that Einstein's theory passes this extraordinarily stringent test with a fractional accuracy better than 0.4% (see Figures 10 and 11). The clock-comparison experiment for PSR 1913 + 16 thus provides direct experimental proof that changes in gravity propagate at the speed of light, thereby creating a dissipative mechanism in an orbiting system. It necessarily follows that gravitational radiation exists and has a quadrupolar nature.

PSR 1534+12 was discovered just three years ago, in a survey by Aleksander Wolszczan (41) that again used the huge Arecibo telescope to good advantage. This pulsar promises eventually to surpass the results now avail-




Fig. 11: Solid curves correspond to Eqs. (5–7) together with the measured values of \dot{P}_b . Their intersection at a single point (within the experimental uncertainty of about 0.35% in \dot{P}_b), establishes the existence of gravitational waves. Dashed curves correspond to the *predicted* values of parameters r and s ; these quantities should become measurable with a modest improvement in data quality.

able from PSR 1913 + 16. It has orbital period $P_b \approx 10.1$ h, eccentricity $e \approx 0.27$, and mass function $f, \approx 0.31 M_\odot$. Moreover, with a stronger signal and narrower pulse than PSR 1913 + 16, its TOAs have considerably smaller measurement uncertainties, around 3 μ s for five-minute observations. Results based on 15 months of data (39) have already produced significant measurements of four PK parameters: $\dot{\omega}$, γ , r , and s . In recent work not yet published, Wolszczan and I have measured the orbital decay rate, \dot{P}_b , and found it to be in accord with general relativity at about the 20% level. In fact, all measured parameters of the PSR 1534+12 system are consistent within general relativity, and it appears that when the full experimental analysis is complete, Einstein's theory will have passed three more very stringent tests under strong-field and radiative conditions.

I do not believe that general relativity necessarily contains the last valid words to be written about the nature of gravity. The theory is not, of course, a quantum theory, and at its most fundamental level the universe appears to

obey quantum-mechanical rules. Nevertheless, our experiments with binary pulsars show that, whatever the precise directions of future theoretical work may be, the correct theory of gravity must make predictions that are asymptotically close to those of general relativity over a vast range of classical circumstances.

ACKNOWLEDGMENTS

Russell Hulse and I have many individuals to thank for their important work, both experimental and theoretical, without which our discovery of PSR 1913 + 16 could not have borne fruit so quickly or so fully. Most notable among these are Roger Blandford, Thibault Damour, Lee Fowler, Peter McCulloch, Joel Weisberg, and the skilled and dedicated technical staff of the Arecibo Observatory.

REFERENCES

1. A. Hewish, S. J. Bell, J. D. H. Pilkington, P. F. Scott, and R. A. Collins. Observation of a rapidly pulsating radio source. *Nature*, 217:709 - 713, 1968.
2. G. R. Huguenin, J. H. Taylor, L. E. Goad, A. Hartai, G. S. F. Orsten, and A. K. Rodman. New pulsating radio source. *Nature*, 219:576, 1968.
3. T. Gold. Rotating neutron stars as the origin of the pulsating radio sources. *Nature*, 218:731-732, 1968.
4. D. H. Staelin and E. C. Reifenstein, III. Pulsating radio sources near the Crab Nebula. *Science*, 162:1481-1483, 1968.
5. M. I. Large, A. E. Vaughan, and B. Y. Mills. A pulsar supernova association. *Nature*, 220:340-341, 1968.
6. D. W. Richards and J. M. Comella. The period of pulsar NP 0532. *Nature*, 222:551-552, 1969.
7. V. Radhakrishnan and D. J. Cooke. Magnetic poles and the polarization structure of pulsar radiation. *Astrophys. Lett.*, 3:225 - 229, 1969.
8. P. Goldreich and W. H. Julian. Pulsar electrodynamics. *Astrophys. J.*, 157:869 - 880, 1969.
9. R. N. Manchester and W. L. Peters. Pulsar parameters from timing observations. *Astrophys. J.*, 173:221-226, 1972.
10. W. R. Burns and B. G. Clark. Pulsar search techniques. *Astron. Astrophys.*, 2:280-287, 1969.
11. R. N. Manchester, J. H. Taylor, and G. R. Huguenin. New and improved parameters for twenty-two pulsars. *Nature Phys. Sci.*, 240:74, 1972.
12. J. H. Taylor. A sensitive method for detecting dispersed radio emission. *Astron. Astrophys. Supp. Ser.*, 15:367, 1974.
13. J. H. Taylor, R. N. Manchester, and A. G. Lyne. Catalog of 558 pulsars. *Astrophys. J. Supp. Ser.*, 88:529 - 568, 1993.
14. F. Camilo. Millisecond pulsar searches. In A. Alpar, editor, *Lives of the Neutron Stars*, (NATO ASI Series), Dordrecht, 1994. Kluwer.
15. J. H. Taylor. A high sensitivity survey to detect new pulsars. Research proposal submitted to the US National Science Foundation, September, 1972.
16. R. A. Hulse and J. H. Taylor. A high sensitivity pulsar survey. *Astrophys. J. (Letters)*, 191:L59--61, 1974.

17. R. A. Hulse and J. H. Taylor. Discovery of a pulsar in a binary system. *Astrophys. J.*, 195:L51 - L53, 1975.
18. R. A. Hulse and J. H. Taylor. A deep sample of new pulsars and their spatial extent in the galaxy. *Astrophys. J. (Letters)*, 201:L55- L59, 1975.
19. R. A. Hulse. The discovery of the binary pulsar. *Les Prix Nobel 1993*, 58-79. The Nobel Foundation, 1994.
20. J. H. Taylor. Millisecond pulsars: Nature's most stable clocks. *Proc. I. E. E. E.*, 79:1054-1062, 1991.
21. J. H. Taylor, R. A. Hulse, L. A. Fowler, G. E. Gullahorn, and J. M. Rankin. Further observations of the binary pulsar PSR 1913 + 16. *Astrophys. J.*, 206:L53 - L58, 1976.
22. P. M. McCulloch, J. H. Taylor, and J. M. Weisberg. Tests of a new dispersion-removing radiometer on binary pulsar PSR 1913+ 16. *Astrophys. J. (Letters)*, 227:L133-1~137, 1979.
23. J. H. Taylor, L. A. Fowler, and P. M. McCulloch. Measurements of general relativistic effects in the binary pulsar PSR 1913 + 16. *Nature*, 277:437, 1979.
24. J. H. Taylor and J. M. Weisberg. A new test of general relativity: Gravitational radiation and the binary pulsar PSR 1913 + 16. *Astrophys. J.*, 253:908-920, 1982.
25. J. H. Taylor and J. M. Weisberg. Further experimental tests of relativistic gravity using the binary pulsar PSR 1913 + 16. *Astrophys. J.*, 345:434 - 450, 1989.
26. D. R. Stinebring, V. M. Kaspi, D. J. Nice, M. F. Ryba, J. H. Taylor, S. E. Thorsett, and T. H. Hankins. A flexible data acquisition system for timing pulsars. *Rev. Sci. Instrum.*, 63:3551-3555, 1992.
27. R. Blandford and S. A. Teukolsky. Arrival-time analysis for a pulsar in a binary system. *Astrophys. J.*, 205:580-591, 1976.
28. R. Epstein. The binary pulsar: Post Newtonian timing effects. *Astrophys. J.*, 216:92-100, 1977.
29. M. P. Haugan. Post-Newtonian arrival-time analysis for a pulsar in a binary system. *Astrophys. J.*, 296:1- 12, 1985.
30. T. Damour and N. Deruelle. General relativistic celestial mechanics of binary systems. I. The post-Newtonian motion. *Ann. Inst. H. Poincare (Physique Thiorique)*, 43:107-132, 1985.
31. T. Damour and N. Deruelle. General relativistic celestial mechanics of binary systems. II. The post-Newtonian timing formula. *Ann. Inst. H. Poincare' (Physique Theorique)*, 44:263 - 292, 1986.
32. T. Damour and J. H. Taylor. Strong-field tests of relativistic gravity and binary pulsars. *Phys. Rev. D*, 45:1840- 1868, 1992.
33. L. A. Rawley, J. H. Taylor, and M. M. Davis. Fundamental astrometry and millisecond pulsars. *Astrophys. J.*, 326:947 - 953, 1988.
34. M. F. Ryba and J. H. Taylor. High precision timing of millisecond pulsars. I. Astrometry and masses of the PSR 1855+09 system. *Astrophys. J.*, 371:739-748, 1991.
35. V. M. Kaspi, J. H. Taylor, and M. Ryba. High-precision timing of millisecond pulsars. III. Long-term monitoring of PSRs B1855+09 and B1937+21. *Astrophys. J.*, in press, 1994.
36. J. H. Taylor and R. J. Dewey. Improved parameters for four binary pulsars. *Astrophys. J.*, 332:770 - 776, 1988.
37. S. E. Thorsett, Z. Arzoumanian, M. M. McKinnon, and J. H. Taylor. The masses of two binary neutron star systems. *Astrophys. J. (Letters)*, 405:L29-L32, 1993.
38. J. H. Taylor, A. Wolszczan, T. Damour, and J. M. Weisberg. Experimental constraints on strong-field relativistic gravity. *Nature*, 355: 132 - 136, 1992.
39. J. H. Taylor. Testing relativistic gravity with binary and millisecond pulsars. In R. J. Gleiser, C. N. Kozameh, and O. M. Moreschi, editors, *General Relativity and Gravitation 1992*, pages 287-294, Bristol, 1993. Institute of Physics Publishing.

40. T. Damour and J. H. Taylor. On the orbital period change of the binary pulsar PSR 1913+16. *Astrophys. J.*, 366:501 -511, 1991.
41. A. Wolszczan. A nearby 37.9 ms radio pulsar in a relativistic binary system. *Nature*, 350:688-690, 1991.